



全口牙結石清除注意事項



若有其他特殊問題，無法自行處理請速到本院或來電詢問。

牙科專線：(08)832-0020、(08)832-9966轉分機3062

一、牙結石清除後2~3天：

對冷的東西會有些敏感，且勿吃刺激食物。(如：熱、冰、辣)



二、牙齦流血：

如果牙齦組織不好，所以術後會稍微流血，是屬正常現象(牙齦組織好者，則不會有此情形) 若仍有流血請回門診處理。

三、醫師建議：

每半年全口牙結石清除一次，並做口腔健康檢查。



安泰醫療社團法人安泰醫院

ANTAI MEDICAL CARE CORPORATION ANTAI TIAN-SHENG MEMORIAL HOSPITAL

牙科 Dentistry





Precautions after full mouth scaling



1. Sensitive to cold food or beverage within 2~3 days after scaling, and do **NOT** eat irritated food (ex: hot, ice, spicy)



2. If the gum has inflammation

there will be some bleeding after scaling (otherwise it seldom happen); However, if the bleeding tendency is high and lasts too long, then OPD check-up as soon as possible is recommended.



3. Routine full mouth scaling and examination every 6 months is recommended.

4. If there is any questions

please call to us or visit us during our working hours.



Tel: (+886) 8-8320020; (+886) 8-8329966 ext. 3062

Add: No.210, Sec. 1, Zhongzheng Rd., Donggang Township, Pingtung County 928, Taiwan (R.O.C.)



安泰醫療社團法人安泰醫院

ANTAI MEDICAL CARE CORPORATION ANTAI TIAN-SHENG MEMORIAL HOSPITAL

牙科 Dentistry

